

Review on Obesity Causes and Prevention

Prachi Sara

All India Institute of Medical Sciences Delhi, India

Email: prachisara@gmail.com

ABSTRACT

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have an adverse effect on health. It is defined by body mass index (BMI) and further evaluated in terms of fat distribution via the waist-hip ratio and total cardiovascular risk factors. Although there are genetic, behavioral, metabolic and hormonal influences on body weight, obesity occurs when you take in more

Keywords: Obesity, causes, prevention.

calories than you burn through exercise and normal daily activities. Your body stores these excess calories as fat. It can be totally prevented through daily exercise, a healthy diet, and a long-term commitment to watch what you eat and drink. Obesity are now dramatically on the rise in low- and middle-income countries, particularly in urban settings.

INTRODUCTION

Obesity is a complex disease involving an excessive amount of body fat. Obesity isn't just a cosmetic concern. It is a medical problem that increases your risk of other diseases and health problems, such as heart disease, diabetes, high blood pressure and certain cancers. There are many reasons why some people have difficulty avoiding obesity. Usually, obesity results from a combination of inherited factors, combined with the environment and personal diet and exercise choices [1] [2]. The good news is that even modest weight loss can improve or prevent the health problems associated with obesity. Dietary changes, increased physical activity and behavior changes can help you lose weight. Prescription medications and weight-loss procedures are additional options for treating obesity. Obesity is diagnosed when your body mass index (BMI) is 30 or higher [3]. To determine your body mass index, divide your weight in pounds by your height in inches squared and multiply by 703. Or divide your weight in kilograms by your height in meters squared. For most people, BMI provides a reasonable estimate of body fat [4]. However, BMI doesn't directly measure

body fat, so some people, such as muscular athletes, may have a BMI in the obesity category even though they don't have excess body fat [5]. Obesity can diminish your overall quality of life. You may not be able to do things you used to do, such as participating in enjoyable activities. You may avoid public places. People with obesity may even encounter discrimination [6].

Causes

Family inheritance and influences: The genes you inherit from your parents may affect the amount of body fat you store, and where that fat is distributed. Genetics may also play a role in how efficiently your body converts food into energy, how your body regulates your appetite and how your body burns calories during exercise. Obesity tends to run in families. That's not just because of the genes they share. Family members also tend to share similar eating and activity habits [7] [8].

Lifestyle choices

- Unhealthy diet. A diet that's high in calories, lacking in fruits and vegetables, full of fast food, and laden with high-calorie beverages and oversized portions contributes to weight gain.

- Liquid calories. People can drink many calories without feeling full, especially calories from alcohol. Other high-calorie beverages, such as sugared soft drinks, can contribute to significant weight gain [9].
- Inactivity. If you have a sedentary lifestyle, you can easily take in more calories every day than you burn through exercise and routine daily activities. Looking at computer, tablet and phone screens is a sedentary activity. The number of hours you spend in front of a screen is highly associated with weight gain.

Certain diseases and medications:

In some people, obesity can be traced to a medical cause, such as Prader-Willi syndrome, Cushing syndrome and other conditions [10]. Medical problems, such as arthritis, also can lead to decreased activity, which may result in weight gain. Some medications can lead to weight gain if you don't compensate through diet or activity. These medications include some antidepressants, anti-seizure medications, diabetes medications, antipsychotic medications, steroids and beta blockers.

Social and economic issues:

Social and economic factors are linked to obesity. Avoiding obesity is difficult if you don't have safe areas to walk or exercise [11]. Similarly, you may not have been taught healthy ways of cooking, or you may not have access to healthier foods. In addition, the people you spend time with may influence your weight — you're more likely to develop obesity if you have friends or relatives with obesity.

Age:

Obesity can occur at any age, even in young children. But as you age, hormonal changes and a less active lifestyle increase your risk of obesity. In addition, the amount of muscle in your body tends to decrease with age [12]. Generally, lower muscle mass leads to a decrease in metabolism. These changes also reduce calorie needs, and can make it harder to keep off excess weight. If you don't consciously control what you eat and

become more physically active as you age, you'll likely gain weight [13].

Other factors:

- Pregnancy: Weight gain is common during pregnancy. Some women find this weight difficult to lose after the baby is born. This weight gain may contribute to the development of obesity in women. Breast-feeding may be the best option to lose the weight gained during pregnancy.
- Quitting smoking: Quitting smoking is often associated with weight gain. And for some, it can lead to enough weight gain to qualify as obesity. Often, this happens as people use food to cope with smoking withdrawal [14]. In the long run, however, quitting smoking is still a greater benefit to your health than is continuing to smoke. Your doctor can help you prevent weight gain after quitting smoking.
- Lack of sleep: Not getting enough sleep or getting too much sleep can cause changes in hormones that increase your appetite. You may also crave foods high in calories and carbohydrates, which can contribute to weight gain [15].
- Stress: Many external factors that affect your mood and well-being may contribute to obesity. People often seek more high-calorie food when experiencing stressful situations.
- Microbiome: Your gut bacteria are affected by what you eat and may contribute to weight gain or difficulty losing weight.
- Previous attempts to lose weight: Previous attempts of weight loss followed by rapid weight regain may contribute to further weight gain. This phenomenon, sometimes called yo-yo dieting, can slow your metabolism [16].

Even if you have one or more of these risk factors, it doesn't mean that you're destined to develop obesity. You can

counteract most risk factors through diet, physical activity and exercise, and behavior changes [16].

Complications

People with obesity are more likely to develop a number of potentially serious health problems, including:

- Heart disease and strokes: Obesity makes you more likely to have high blood pressure and abnormal cholesterol levels, which are risk factors for heart disease and strokes [17].
- Type 2 diabetes: Obesity can affect the way your body uses insulin to control blood sugar levels. This raises your risk of insulin resistance and diabetes.
- Certain cancers: Obesity may increase your risk of cancer of the uterus, cervix, endometrium, ovary, breast, colon, rectum, esophagus, liver, gallbladder, pancreas, kidney and prostate [18].
- Digestive problems: Obesity increases the likelihood that you'll develop heartburn, gallbladder disease and liver problems.
- Gynecological and sexual problems: Obesity may cause infertility and irregular periods in women. Obesity also can cause erectile dysfunction in men.
- Sleep apnea: People with obesity are more likely to have sleep apnea, a potentially serious disorder in which breathing repeatedly stops and starts during sleep.
- Osteoarthritis: Obesity increases the stress placed on weight-bearing joints, in addition to promoting inflammation within the body. These factors may lead to complications such as osteoarthritis.

Prevention

Whether you're at risk of obesity, currently overweight or at a healthy weight, you can take steps to prevent unhealthy weight gain and related health problems. Not surprisingly, the steps to prevent weight gain are the same as the

steps to lose weight: daily exercise, a healthy diet, and a long-term commitment to watch what you eat and drink [19].

- Exercise regularly. You need to get 150 to 300 minutes of moderate-intensity activity a week to prevent weight gain. Moderately intense physical activities include fast walking and swimming.
- Follow a healthy-eating plan. Focus on low-calorie, nutrient-dense foods, such as fruits, vegetables and whole grains. Avoid saturated fat and limit sweets and alcohol. Eat three regular meals a day with limited snacking. You can still enjoy small amounts of high-fat, high-calorie foods as an infrequent treat. Just be sure to choose foods that promote a healthy weight and good health most of the time.
- Know and avoid the food traps that cause you to eat. Identify situations that trigger out-of-control eating. Try keeping a journal and write down what you eat, how much you eat, when you eat, how you're feeling and how hungry you are. After a while, you should see patterns emerge. You can plan ahead and develop strategies for handling these types of situations and stay in control of your eating behaviors [20].
- Monitor your weight regularly. People who weigh themselves at least once a week are more successful in keeping off excess pounds. Monitoring your weight can tell you whether your efforts are working and can help you detect small weight gains before they become big problems.
- Be consistent. Sticking to your healthy-weight plan during the week, on the weekends, and amidst vacation and holidays as much as possible increases your chances of long-term success.

CONCLUSION

Obesity is a choice. It takes work and effort to stay in shape, but it can be fun. You must eat healthy, stay active, and make good life choices. But healthy foods can be delicious, there are plenty of activities to do that are so much fun, and making good life choices can be rewarding in multiple ways. To help cut obesity rates, people need to understand

the effects of eating at the fast food restaurants, eating the entire calorie packed foods, and the importance of encouraging healthy lifestyles for children [11]. There are many problems in world today, but we can remove one from the list by helping to cut obesity rates in modern world.

REFERENCES

1. Ackroyd R, Mouiel J, Chevallier J M, Daoud F. Cost-effectiveness and budget impact of obesity surgery in patients with type-2 diabetes in three European countries. 2006.
2. Agras WS, Berkowitz RI, Arnow BA, Telch CF, Marnell M, Henderson J, et al. Maintenance following a very-low-calorie diet. 1996.
3. Anselmino M, Bammer T, Cebrian JMF, Daoud F, Romagnoli G, Torres A. Cost-effectiveness and budget impact of obesity surgery in patients with type 2 diabetes in three European countries (II). *Obesity Surgery*. 2009;19(11):1542-1549. [PubMed]
4. Arai K, Miura J, Ohno M, Yokoyama J, Ikeda Y. Comparison of clinical usefulness of very-low-calorie diet and supplemental low-calorie diet.
5. Borg P, Kukkonen-Harjula K, Fogelholm M, Pasanen M. Effects of walking or resistance training on weight loss maintenance in obese, middle-aged men: a randomized trial. *International Journal of Obesity and Related Metabolic Disorders*. 2002.
6. British Association of Plastic Reconstructive and Aesthetic Surgeons and Royal College of Surgeons. Commissioning guide: massive weight loss body contouring. London: Royal College of Surgeons; 2014.
7. Buse JB, Caprio S, Cefalu WT, Ceriello A, Del PS, Inzucchi SE, et al. How do we define cure of diabetes? *Diabetes Care*. 2009;
8. Christensen P, Bliddal H, Riecke BF, Leeds AR, Astrup A, Christensen R. Blood pressure and lipids improvement after very low energy diet (VLED) or low energy diet (LED) in obese patients with knee osteoarthritis: A randomised controlled trial. *Atherosclerosis*. 2009.
9. Christensen P, Bliddal H, Riecke BF, Leeds AR, Astrup A, Christensen R. Comparison of a low-energy diet and a very low-energy diet in sedentary obese individuals: A pragmatic randomized controlled trial. 2011.
10. Counterweight Project Team. Influence of body mass index on prescribing costs and potential cost savings of a weight management programme in primary care. *Journal of Health Services Research and Policy*. 2008.
11. Cremieux PY, Eapen S, Trask SW, Ghosh A. Weighing the clinical benefits and economic impact of bariatric surgery in morbidly obese patients with diabetes. *Canadian Journal of Diabetes*. 2011.
12. Curtis L. Unit costs of health and social care 2013. Canterbury: Personal Social Services Research Unit, University of Kent; 2013.
13. Delbridge EA, Prendergast LA, Pritchard JE, Proietto J. One-year weight maintenance after significant weight loss in healthy overweight and obese subjects: does diet composition matter? *American Journal of Clinical Nutrition*. 2009.
14. Dixon JB, Zimmet P, Alberti KG, Rubino F. Bariatric surgery: an IDF

- statement for obese Type 2 diabetes. *Diabetic Medicine*. 2011.
15. Dixon S, Currie CJ, McEwan P. Utility values for obesity and preliminary analysis of the Health Outcomes Data Repository.
 16. Finkelstein EA, Allaire BT, DiBonaventura MD, Burgess SM. Direct and indirect costs and potential cost savings of laparoscopic adjustable gastric banding among obese patients with diabetes. *Journal of Occupational and Environmental Medicine*. 2011.
 17. Fogelholm M, Kukkonen-Harjula K, Nenonen A, Pasanen M. Effects of walking training on weight maintenance after a very-low-energy diet in premenopausal obese women: a randomized controlled trial.
 18. Gebhard RL, Prigge WF, Ansel HJ, Schlasner L, Ketover SR, Sande D, et al. The role of gallbladder emptying in gallstone formation during diet-induced rapid weight loss. *Hepatology*. 1996;
 19. Hoerger TJ, Zhang P, Segel JE, Kahn HS, Barker LE, Couper S. Cost-effectiveness of bariatric surgery for severely obese adults with diabetes.
 20. Hunger M, Schunk M, Meisinger C, Peters A, Holle R. Estimation of the relationship between body mass index and EQ-5D health utilities in individuals with type 2 diabetes: evidence from the population-based KORA studies.